

Climate sensitivity across the RCEMIP simulations




Tobias Becker

Max Planck Institute for Meteorology




RCEMIP provides a wide spectrum of models of different complexity


CRM & LES
(small domain)



CRM
(large domain)



GCM



- simulation length:
 - 1000 days (GCM)
 - 100 days (CRM)
 - 50 days (LES)
- three prescribed SSTs:
295 K, 300 K, 305 K
- details on RCE setup in
Wing et al., 2018

Estimating climate sensitivity from simulations with prescribed surface temperature

$$dT = \frac{dR_{\text{TOA}}}{\lambda}$$

$$\text{ECS} = \frac{F_{2\times\text{CO}_2}}{\lambda}$$

Estimating climate sensitivity from simulations with prescribed surface temperature

$$dT = \frac{dR_{\text{TOA}}}{\lambda} \quad \Longleftrightarrow \quad \lambda = \frac{dR_{\text{TOA}}}{dT}$$

$$\text{ECS} = \frac{F_{2\times\text{CO}_2}}{\lambda}$$

Estimating climate sensitivity from simulations with prescribed surface temperature

$$dT = \frac{dR_{\text{TOA}}}{\lambda} \quad \Longleftrightarrow \quad \lambda = \frac{dR_{\text{TOA}}}{dT} = \frac{dR_{\text{TOA},\text{lw}}}{dT} + \frac{dR_{\text{TOA},\text{sw}}}{dT}$$

$$\text{ECS} = \frac{F_{2\times\text{CO}_2}}{\lambda}$$

Estimating climate sensitivity from simulations with prescribed surface temperature

$$dT = \frac{dR_{\text{TOA}}}{\lambda} \quad \Longleftrightarrow \quad \lambda = \frac{dR_{\text{TOA}}}{dT} = \frac{dR_{\text{TOA},\text{lw}}}{dT} + \frac{dR_{\text{TOA},\text{sw}}}{dT}$$

$$\frac{dR_{\text{TOA},\text{clr}}}{dT} = \frac{dR_{\text{TOA},\text{clr},\text{lw}}}{dT} + \frac{dR_{\text{TOA},\text{clr},\text{sw}}}{dT}$$

$$\text{ECS} = \frac{F_{2\times\text{CO}_2}}{\lambda}$$

Estimating climate sensitivity from simulations with prescribed surface temperature

$$dT = \frac{dR_{\text{TOA}}}{\lambda} \quad \Longleftrightarrow \quad \lambda = \frac{dR_{\text{TOA}}}{dT} = \frac{dR_{\text{TOA},\text{lw}}}{dT} + \frac{dR_{\text{TOA},\text{sw}}}{dT}$$

$$\frac{dR_{\text{TOA},\text{clr}}}{dT} = \frac{dR_{\text{TOA},\text{clr},\text{lw}}}{dT} + \frac{dR_{\text{TOA},\text{clr},\text{sw}}}{dT}$$

$$\text{ECS} = \frac{F_{2\times\text{CO}_2}}{\lambda} \quad \Longleftrightarrow \quad \text{ECS} = \frac{F_{2\times\text{CO}_2}}{\frac{dR_{\text{TOA}}}{dT}}$$

Estimating climate sensitivity from simulations with prescribed surface temperature

$$dT = \frac{dR_{\text{TOA}}}{\lambda} \quad \Longleftrightarrow \quad \lambda = \frac{dR_{\text{TOA}}}{dT} = \frac{dR_{\text{TOA},\text{lw}}}{dT} + \frac{dR_{\text{TOA},\text{sw}}}{dT}$$

$$\frac{dR_{\text{TOA},\text{clr}}}{dT} = \frac{dR_{\text{TOA},\text{clr},\text{lw}}}{dT} + \frac{dR_{\text{TOA},\text{clr},\text{sw}}}{dT}$$

$$\text{ECS} = \frac{F_{2\times\text{CO}_2}}{\lambda} \quad \Longleftrightarrow \quad \text{ECS} = \frac{F_{2\times\text{CO}_2}}{\frac{dR_{\text{TOA}}}{dT}} \rightarrow 3.7 \text{ W m}^{-2}$$

Estimating climate sensitivity from simulations with prescribed surface temperature

$$dT = \frac{dR_{\text{TOA}}}{\lambda} \iff \lambda = \frac{dR_{\text{TOA}}}{dT} = \frac{dR_{\text{TOA},\text{lw}}}{dT} + \frac{dR_{\text{TOA},\text{sw}}}{dT}$$
$$\frac{dR_{\text{TOA},\text{clr}}}{dT} = \frac{dR_{\text{TOA},\text{clr},\text{lw}}}{dT} + \frac{dR_{\text{TOA},\text{clr},\text{sw}}}{dT}$$

$$\text{ECS} = \frac{F_{2\times\text{CO}_2}}{\lambda} \iff \text{ECS} = \frac{F_{2\times\text{CO}_2}}{\frac{dR_{\text{TOA}}}{dT}} \rightarrow 3.7 \text{ W m}^{-2}$$

two ECS estimates per model: 305 K - 300 K & 300 K - 295 K
(skipping the first 50 days for GCM / CRM, and the first 25 days for LES)

Climate sensitivity across different RCEMIP model setups

avg


1.30
1.0
1.30
1.0

1.29
1.0
1.29
1.0


1.0
1.0
1.29
1.0

small domain:

- results independent of resolution
- climate sensitivity increases with temperature (esp. clear sky)




Climate sensitivity across different RCEMIP model setups



small domain:

- results independent of resolution
- climate sensitivity increases with temperature (esp. clear sky)
- high climate sensitivity (RCE!), but within CMIP5 model range

Climate sensitivity across different RCEMIP model setups




small domain:

- results independent of resolution
- climate sensitivity increases with temperature (esp. clear sky)
- high climate sensitivity (RCE!), but within CMIP5 model range

large domain:

- standard deviation much higher (outside of CMIP5 model range)
- even negative values possible (infinite climate sensitivity!)

Climate sensitivity across different RCEMIP model setups



small domain:

- results independent of resolution
- climate sensitivity increases with temperature (esp. clear sky)
- high climate sensitivity (RCE!), but within CMIP5 model range


large domain:

- standard deviation much higher (outside of CMIP5 model range)
- even negative values possible (infinite climate sensitivity!)

GCMs:

- spread even larger
- smaller mean climate sensitivity
- smaller spread for clear-sky fluxes

Climate sensitivity across different RCEMIP model setups



small domain:

- results independent of resolution
- climate sensitivity increases with temperature (esp. clear sky)
- high climate sensitivity (RCE!), but within CMIP5 model range

large domain:


- standard deviation much higher (outside of CMIP5 model range)
- even negative values possible (infinite climate sensitivity!)

GCMs:


- spread even larger
- smaller mean climate sensitivity
- smaller spread for clear-sky fluxes

UCP Berlin
02/2019

MPI large domain models: extreme spread in climate sensitivity related to temperature dependence of convective self-aggregation





credits: Hauke Schulz; Tompkins and Semie, 2017




Can we also relate the extreme spread in climate sensitivity across all RCEMIP models to the temperature dependence of convective self-aggregation?



Can we also relate the extreme spread in climate sensitivity across all RCEMIP models to the temperature dependence of convective self-aggregation?




Can we also relate the extreme spread in climate sensitivity across all RCEMIP models to the temperature dependence of convective self-aggregation?




- significant correlation both for GCMs and CRMs
- smaller climate sensitivity in GCMs in line with positive I_{org} / dT

Can we also relate the extreme spread in climate sensitivity across all RCEMIP models to the temperature dependence of convective self-aggregation?




- significant correlation both for GCMs and CRMs
- smaller climate sensitivity in GCMs in line with positive I_{org} / dT
- a necessary condition for extreme climate sensitivities seems to be the according tendency in self-aggregation

Can we also relate the extreme spread in climate sensitivity across all RCEMIP models to the temperature dependence of convective self-aggregation?





- significant correlation both for GCMs and CRMs
- smaller climate sensitivity in GCMs in line with positive I_{org} / dT
- a necessary condition for extreme climate sensitivities seems to be the according tendency in self-aggregation
- clear-sky TOA fluxes still show the same correlation
→ mechanism is independent of clouds

Can we also relate the extreme spread in climate sensitivity across all RCEMIP models to the temperature dependence of convective self-aggregation?




- significant correlation both for GCMs and CRMs
- smaller climate sensitivity in GCMs in line with positive I_{org} / dT
- a necessary condition for extreme climate sensitivities seems to be the according tendency in self-aggregation
- clear-sky TOA fluxes still show the same correlation
→ mechanism is independent of clouds
- changes in the clear-sky radiation budget only depend on longwave radiation
→ mechanism: convective self-aggregation leads to a stronger overturning circulation, stronger drying in the subsidence regions and increased outgoing longwave radiation


GCMs and one CRM (UCLA-LES) also show a strong dependency on shallow cloud fraction in the subsidence regime
($R_{TOA,lw} > 250 \text{ W m}^{-2}$; $R_{TOA,sw,up} > 100 \text{ W m}^{-2}$)



GCMs and one CRM (UCLA-LES) also show a strong dependency on shallow cloud fraction in the subsidence regime
($R_{TOA,lw} > 250 \text{ W m}^{-2}$; $R_{TOA,sw,up} > 100 \text{ W m}^{-2}$)




GCMs and one CRM (UCLA-LES) also show a strong dependency on shallow cloud fraction in the subsidence regime
($R_{TOA, lw} > 250 \text{ W m}^{-2}$; $R_{TOA, sw, up} > 100 \text{ W m}^{-2}$)




- for GCMs: significantly higher correlation of dR_{TOA} / dT with f_{sc} / dT than with I_{org} / dT
- clear-sky fluxes do not show any correlation

GCMs and one CRM (UCLA-LES) also show a strong dependency on shallow cloud fraction in the subsidence regime
($R_{TOA,lw} > 250 \text{ W m}^{-2}$; $R_{TOA,sw,up} > 100 \text{ W m}^{-2}$)



- for GCMs: significantly higher correlation of dR_{TOA} / dT with f_{sc} / dT than with I_{org} / dT
- clear-sky fluxes do not show any correlation
- longwave fluxes do not show any correlation

GCMs and one CRM (UCLA-LES) also show a strong dependency on shallow cloud fraction in the subsidence regime
($R_{TOA,lw} > 250 \text{ W m}^{-2}$; $R_{TOA,sw,up} > 100 \text{ W m}^{-2}$)



- for GCMs: significantly higher correlation of dR_{TOA} / dT with f_{sc} / dT than with l_{org} / dT
- clear-sky fluxes do not show any correlation
- longwave fluxes do not show any correlation
- changes of shallow clouds in the subsidence region with T strongly affect climate sensitivity via their influence on how much shortwave radiation they reflect back to space

Summary

- On the small domain (in the absence of convective self-aggregation), climate sensitivity estimates are
 - robust (similar for CRMs & LES models)
 - in the same range as in CMIP5
 - increasing with temperature




Summary

- On the small domain (in the absence of convective self-aggregation), climate sensitivity estimates are
 - robust (similar for CRMs & LES models)
 - in the same range as in CMIP5
 - increasing with temperature
- On the large domain and in GCMs, differences in convective organization and clouds can lead to very different climate sensitivities across the RCEMIP simulations


Summary

- On the small domain (in the absence of convective self-aggregation), climate sensitivity estimates are
 - robust (similar for CRMs & LES models)
 - in the same range as in CMIP5
 - increasing with temperature
- On the large domain and in GCMs, differences in convective organization and clouds can lead to very different climate sensitivities across the RCEMIP simulations
- Two mechanisms are responsible for extreme climate sensitivities:
 - increased convective self-aggregation leads to a stronger overturning circulation, stronger drying in the subsidence regions and increased outgoing longwave radiation




Summary

- On the small domain (in the absence of convective self-aggregation), climate sensitivity estimates are
 - robust (similar for CRMs & LES models)
 - in the same range as in CMIP5
 - increasing with temperature
- On the large domain and in GCMs, differences in convective organization and clouds can lead to very different climate sensitivities across the RCEMIP simulations
- Two mechanisms are responsible for extreme climate sensitivities:
 - increased convective self-aggregation leads to a stronger overturning circulation, stronger drying in the subsidence regions and increased outgoing longwave radiation
 - more shallow clouds in the subsidence region reflect more shortwave radiation (this mechanism is mainly important in GCMs!)




Summary

- On the small domain (in the absence of convective self-aggregation), climate sensitivity estimates are
 - robust (similar for CRMs & LES models)
 - in the same range as in CMIP5
 - increasing with temperature
- On the large domain and in GCMs, differences in convective organization and clouds can lead to very different climate sensitivities across the RCEMIP simulations
- Two mechanisms are responsible for extreme climate sensitivities:
 - increased convective self-aggregation leads to a stronger overturning circulation, stronger drying in the subsidence regions and increased outgoing longwave radiation
 - more shallow clouds in the subsidence region reflect more shortwave radiation (this mechanism is mainly important in GCMs!)
- Climate sensitivity is overall larger in CRMs than in GCMs



Summary

- On the small domain (in the absence of convective self-aggregation), climate sensitivity estimates are
 - robust (similar for CRMs & LES models)
 - in the same range as in CMIP5
 - increasing with temperature
- On the large domain and in GCMs, differences in convective organization and clouds can lead to very different climate sensitivities across the RCEMIP simulations
- Two mechanisms are responsible for extreme climate sensitivities:
 - increased convective self-aggregation leads to a stronger overturning circulation, stronger drying in the subsidence regions and increased outgoing longwave radiation
 - more shallow clouds in the subsidence region reflect more shortwave radiation (this mechanism is mainly important in GCMs!)
- Climate sensitivity is overall larger in CRMs than in GCMs
- Open Question: Why do the different RCEMIP models handle the mechanisms that lead to extreme climate sensitivities so differently?



Summary

- On the small domain (in the absence of convective self-aggregation), climate sensitivity estimates are
 - robust (similar for CRMs & LES models)
 - in the same range as in CMIP5
 - increasing with temperature
- On the large domain and in GCMs, differences in convective organization and clouds can lead to very different climate sensitivities across the RCEMIP simulations
- Two mechanisms are responsible for extreme climate sensitivities:
 - increased convective self-aggregation leads to a stronger overturning circulation, stronger drying in the subsidence regions and increased outgoing longwave radiation
 - more shallow clouds in the subsidence region reflect more shortwave radiation (this mechanism is mainly important in GCMs!)
- Climate sensitivity is overall larger in CRMs than in GCMs
- Open Question: Why do the different RCEMIP models handle the mechanisms that lead to extreme climate sensitivities so differently?

